

# Newts under siege: range expansion of *Triturus pygmaeus* isolates populations of its sister species

G. Espregueira Themudo<sup>1\*</sup> and J. W. Arntzen<sup>2</sup>

<sup>1</sup>CIBIO, Centro de Investigação em

Biodiversidade e Recursos Genéticos, Campus Agrário de Vairão, 4485-661 Vairão, Portugal,

<sup>2</sup>National Museum of Natural History, PO Box 9517, 2300 RA Leiden, the Netherlands

\*Correspondence: G. Espregueira Themudo, CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, Campus Agrário de Vairão, 4485-661 Vairão, Portugal. E-mail: themudo@mail.icav.up.pt

## ABSTRACT

The newt species *Triturus marmoratus* and *Triturus pygmaeus* are both present in central Portugal where they have parapatric distributions. We used four genetic markers to determine which species was present in 31 populations. In the centre of the study area we found a *T. marmoratus* enclave. Despite small interpopulation distances, hybridization is locally rare. We built several models to try to explain this distribution using environmental data. The best model, chosen by Akaike's Information Criterion, relates the presence of *T. marmoratus* with the temperature in July, the relief of the landscape, and a higher use of the land for orchards. The current distribution can best be explained by *T. pygmaeus* expanding north and replacing *T. marmoratus*, the latter only persisting where ecological conditions are relatively favourable.

## Keywords


Akaike's Information Criterion, allozymes, Amphibia, enclaves, mosaic hybrid zones, parapatry.

## INTRODUCTION

Secondary contact between closely related species often generates spectacular events and provides insight into the evolutionary process. Hybrid zones formed this way are considered 'natural laboratories' or 'windows on evolutionary processes' (Hewitt, 1988; Harrison, 1990). Evolution is not a static process and to understand it, we must look into dynamic systems like these. The dynamics of species ranges are not independent of ecological conditions, competition with sister species, or dispersal capabilities. Present-day distributions are echoes from past events. One particular event that provides clues to the relative movement of species is the formation of enclaves. Enclaves are here defined as populations of one species *completely* surrounded by populations of closely related species and genetically isolated from other populations of the same species (Arntzen, 1978; in geographical terms, they are simultaneously exclaves and enclaves). This is reminiscent of the 'internal parapatry' concept of Key (1981) that, however, does not deal with disjunct distributions. Mosaic hybrid zones are bimodal hybrid zones with few hybrids and predominantly parental genotypes present. The contact between the (sub)species is more strongly shaped by ecological constraints than by genetic interactions. In a recent review, Jiggins & Mallet (2000) go one step further and suggest that ecology contributes more to speciation than genetic incompatibility. Well-known mosaic hybrid zones are for example those in *Gryllus* crickets (Rand & Harrison, 1989), *Chorthippus* grasshoppers (Bridle *et al.*, 2001), and *Mytilus* mussels (Bierne *et al.*, 2003). Examples of mosaic hybrid zones in salamanders are *Triturus cristatus* and

*Triturus marmoratus* in western France (Arntzen & Wallis, 1991) and *Plethodon cinereus* and *Plethodon shenandoah* in the Appalachian Mountains of North America (Jaeger, 1970, 1971; see also Sites *et al.*, 2001). Although they present patches of populations of one species distributed among patches of the other, these are not all necessarily enclaves because dispersal among patches may be frequent. To the best of our knowledge the only enclaves recorded in the literature are those for *Bombina* toads in central Europe (Arntzen, 1978) and *Triturus* newts in western France and the northern Balkans (Arntzen & Wallis, 1991, 1999). Perhaps enclaves are more likely to arise in organisms with structured populations and low dispersal capability than in organisms that disperse well. Amphibian populations in particular are well delimited because of their dependence on water for reproduction and they have low individual mobility. Once formed, enclaves will take some time to dissolve, or be stable or disappear by reconnection to the main distribution.

The two species of marbled newts living in the Iberian Peninsula, *T. marmoratus* (Latreille, 1800) and *Triturus pygmaeus* (Wolterstorff, 1905), have a parapatric distribution. Some reports, however, indicated the presence of *T. marmoratus* where only *T. pygmaeus* was expected, near Caldas da Rainha and a spatial-environmental model for the two species, suggests that the local conditions may indeed be favourable to *T. marmoratus* (see the southernmost record in Fig. 1c in Arntzen, 2006). This would indicate an area of sympatry or a mosaic distribution. A mosaic distribution would point to differential ecological requirements with patches where the environmental conditions are more suitable for one species than for the other. An intuitive



**Figure 1** Proportion of membership of individuals to species 1 (= *Triturus pygmaeus*) using the software STRUCTURE 2.1 (Pritchard *et al.*, 2000). Data are separated out for samples with 0, 25, 50 and 75% missing data to show that, in practical terms, discriminatory power is independent of the *loci* used.

explanation from fieldwork across the Iberian Peninsula is that *T. pygmaeus* thrives in ephemeral ponds with a fluctuating reproductive output and that *T. marmoratus* thrives in smaller, more permanent water bodies (e.g. springs) with regular but low annual recruitment reaching metamorphosis (J. W. Arntzen, unpublished data). To confirm the record at Caldas da Rainha and to learn more about the events that lead to this occurrence and its extent, we conducted a detailed study. Adult marbled newts have clearly distinguishable morphologies while embryos and larvae are difficult to identify. Conversely, embryos and larvae are easy to find at the aftermath of the reproductive season while adults may be elusive. To facilitate a fast, detailed and reliable surveying, we employed genetic markers for species identification (Espregueira Themudo & Arntzen, 2007). In the present paper, we looked into the spatial structure of the distribution of the two species of marbled newts and determine if and what ecological constraints are shaping it. We also analysed the presence of hybrids in our sample.

## METHODS

### Sampling

In April 2003, we searched around Caldas da Rainha, in central Portugal, for water bodies containing marbled newts. The region is characterized by flat dunes on the coast and an undulating agricultural land, with an abundance of orchards in the centre. We located and sampled 31 populations of *T. marmoratus* and *T. pygmaeus*, over 1600 km<sup>2</sup>. Breeding female newts obtain one or several spermatophores from which eggs are internally fertilized and then deposited one by one, mostly on aquatic vegetation. The process takes place over a *c.* 2-month period. To increase random collection and to reduce pseudoreplication, we screened

the entire accessible area of each pond and collected no more than one egg per leaf or two eggs per plant. In marbled newts adult population size can be very small, especially in small water bodies such as springs (Schoorl & Zuiderwijk, 1981; Jehle *et al.*, 2001, 2005), and this explains why sample size is small on some occasions ( $N < 5$  in four populations). Adults and larvae were captured by dip-netting. Tail tips were collected from adults and larvae were sacrificed. All samples were immediately stored in liquid nitrogen and later transferred to  $-80^{\circ}\text{C}$  until the day they were analysed.

### Genetic data

All the tissue samples (from  $n = 10$  individuals for most locales) were analysed for four allozyme loci: peptidase A (*Pep-A*), peptidase B (*Pep-B*), peptidase D (*Pep-D*), and lactate dehydrogenase (*Ldh-2*; this locus is not yet expressed in embryos), that yield a species-specific enzyme profile using standard starch gel electrophoresis and isoelectric focusing. The genetic signature is consistent with morphological identification of adult marbled newts (Espregueira Themudo & Arntzen, 2007). With the program ARLEQUIN (version 3.1; Excoffier *et al.*, 2005), we tested for departures from Hardy–Weinberg expectations and linkage disequilibrium. We used the program FSTAT (version 2.9.3.2; Goudet, 1995) to calculate *F*-statistics (Weir & Cockerham, 1984) and, to detect population substructuring, we analysed the results with STRUCTURE 2.1 (Pritchard *et al.*, 2000). Using a Markov chain Monte Carlo (MCMC) algorithm, STRUCTURE assigns individuals to a population, or jointly to two or more populations, if their genotypes indicate that they are admixed. This is done assuming a model with *K* populations (where *K* may be unknown), where each is characterized by a set of alleles for each locus. Individuals are assigned to populations as to maximize Hardy–Weinberg and linkage equilibrium. We choose for the ‘admixture model’ because neighbouring populations may interbreed and varied *K* from 1 to 5 with 10,000 generations as the length of burn-in period and 100,000 MCMC generations after burn-in. Convergence occurred before 10,000 in test runs. The value of  $\lambda$  was inferred by the program. We accepted *K* as indicating the true number of genetic partitions when the difference in likelihood between two consecutive values of *K* was maximal.

We investigated the existence of hybrids in our sample by performing a Bayesian model-based clustering as implemented in the software NEWHYBRIDS (Anderson & Thompson, 2002). This computes the posterior probability that each individual belongs to each of six predetermined classes (pure species A, pure species B,  $F_1$ -hybrid,  $F_2$ -hybrid, backcross to species A, and backcross to species B).

### Ecological data

For the spatial environmental analysis in a Geographical Information System (GIS), we selected 21 ecological parameters following Teixeira *et al.* (2001) and an additional explanatory variable (land surface occupied by orchards, arcsin transformed percentages — ORCH) that appeared locally informative. We

used ORCH as a proxy to one or more unidentified variables that — possibly more directly than ORCH — would help to explain the newt distribution. The advantage of the parameter ORCH is that it manifests itself from field observations and that blanket data are readily available (INE, 1999). For all variables, information was available in digital format for Portugal (DGA, 1995). A vegetation map (normalized difference vegetation index or NDVI) was obtained courtesy of the Royal Dutch Meteorological Institute (KNMI). An altitude map was taken from the internet (<http://edcwww.cr.usgs.gov/doc/edchome/datasets/edcdata.html>) and used to produce a relief map by a set of filter operations (Ilwis 2.1, 1997). Maps on the mean January and July temperature were digitalized from the Portuguese climate atlas (SCN, 1974). A hierarchical clustering based on Spearman's correlation coefficient was used to evaluate the level at which ecological information appeared redundant. One variable was selected arbitrarily out of a set of variables that correlated at Spearman's  $r_s > 0.8$ . Sixteen variables were retained: acidity of the soil (ACID), altitude (ALTI), chlorates content of subterranean water (CHLO), frost days (FROD), frost months (FROM), hardness of subterranean water (HARD), humidity of the air (HUMI), insolation (INSO), lithology (LITH), vegetation index (NDVI), orchard land coverage (ORCH), mean annual precipitation (PRET), relief (RELI), the sulphate content of subterranean water (SULP), mean annual temperature (TEMP), and mean July temperature (TJUL). To increase the comparability of their effects, all continuous variables were standardized to an average of zero and a standard deviation of one. The variables were introduced into the GIS analytical software as raster layers with 1 km spatial resolution. Values for 10\*10 km UTM grids were obtained by averaging the data (modal values for the categorical variable LITH).

### Model selection

We used an information-theoretical model selection approach for the statistical analysis of the data (Burnham & Anderson, 2002). Using our field knowledge, we built several *a priori* models that would explain the current distribution of the two species. By contrasting the presence of one species against the other, we circumvented the inclusion of false absence data. To understand the effect of the variable ORCH, we used our available data on fruit growing in the region as dependent variable, and applied a step-wise multiple regression (SMR) with the same environmental data (all but ORCH) as explanatory variables, using the software SPSS version 14 (SPSS Inc., 2005). We then substituted ORCH by the model derived from this analysis in our *a priori* models, and added them to the list as *a posteriori* models. In a second step, we used the small-sample Akaike's Information Criterion (AICc; Burnham & Anderson, 2002) to rank the models and chose the best one. AICc is defined as:

$$\text{AICc} = -2\ln L + 2K + \frac{2K(K+1)}{n-K-1}$$

where  $\ln L$  is the natural logarithm of the likelihood function,  $K$  is the number of parameters from the model, and  $n$  is the sample

**Table 1** Log-likelihood and standard deviation of the number of partitions ( $K$ ) in our data set.  $K$  varied from 1 to 5. The increase in likelihood is not significant other than at  $K = 2$ .

| $K$ | $\log L$ | SD   |
|-----|----------|------|
| 1   | -1737.7  | 5.3  |
| 2   | -961.9   | 0.6  |
| 3   | -887.5   | 1.4  |
| 4   | -849.4   | 30.7 |
| 5   | -802.2   | 6.9  |

size. AIC prevents overfitting the model, by taking the number of parameters into consideration. To fit the models to the data and obtain the log-likelihoods of the models, we used the logistic regression procedure available in SPSS.

### RESULTS

We analysed 398 individuals (101 adults, 50 larvae, and 247 embryos; see allele frequencies in Table S1 in Supplementary Material).


F-statistics shows a high degree of population substructuring ( $F_{st}$ ) in *Pep-A* (0.84) and *Pep-D* (0.66), intermediate in *Ldh-2* (0.41), and low in *Pep-B* (0.14). The exact test on Hardy-Weinberg expectations showed that population 4 presents a significant departure at one locus (*Pep-D*) under Bonferroni correction. Linkage disequilibrium tests are significant at  $P < 0.05$  in two *T. marmoratus* populations that are neighbouring *T. pygmaeus* populations (populations 23 and 31; see Fig. 1 and Table S2 in Supplementary Material).

The model-based clustering method implemented in STRUCTURE 2.1 showed that the most likely number of partitions for the present data is  $K = 2$  (Table 1; for details see Table S2). The partitions correspond to *T. marmoratus* and *T. pygmaeus*. Since the likelihood continues to increase with  $K$ , we looked into the population structure when  $K = 3$ , to examine the possibility that additional substructuring existed. The only result was that one of the clusters split into two equal parts.  $F_{st}$  values are consistent with the relative diagnostic power of the loci (high for *Pep-A* and *Pep-D*, intermediate for *Ldh-2*, and low for *Pep-B*). Accurate species identification is independent of the number of loci studied (Fig. 1).

The spatial distribution of the two groups indicates the existence of a set of *T. marmoratus* populations surrounded by *T. pygmaeus* populations, forming an enclave in the centre of our study area (Fig. 2; note that the isolated pocket is — in proper terms — an exclave of *T. marmoratus* and an enclave of *T. pygmaeus*). Only population 18 had individuals of both species.

The Bayesian-based assignment of individuals to hybrid classes using NEWHYBRIDS indicates that hybridization, backcrossing, and introgression are locally rare. Thirty out of 31 populations were classified as either *T. marmoratus* or *T. pygmaeus* and not both. Population 18 contained both species as well as one individual with about equal probability of being a pure *T. marmoratus* ( $P = 0.53$ ) as having mixed species parentage

**Figure 2** Distribution of marbled newts in Portugal with *Triturus marmoratus* (hatched) and *Triturus pygmaeus* (shaded). The box shows the research area around Caldas da Rainha (a). The detailed map (b) shows the localities with *T. marmoratus* (solid dots) and *T. pygmaeus* (open dots). See Table 1 for population details. Voronoi polygons are used to estimate the contiguous species distribution over the area. Note that the scale of extrapolation at the exterior of the study area is set at c. 6 km as to match the level of interpolation.



[ $P=0.47$ , being either an  $F_2$ -hybrid ( $P=0.35$ ) or a backcross hybrid in direction of *T. marmoratus* ( $P=0.12$ )]. This mixed population (population in grey in Fig. 2) was excluded from the spatial-environmental analysis.

GIS-model 9 (ORCH + TJUL + RELI; see Table 2) showed the lowest AICc value. This model explains 50% of the total variance observed. The presence of *T. marmoratus* relates to the higher abundance of orchards, lower July temperature, and smoother relief than for *T. pygmaeus*. In the subsequent analysis, five environmental variables: ALTI, ACID, HUMI, SULP, and TEMP explained the variable ORCH. This model explained 69% of the variance of ORCH. When, however, ORCH was replaced by this set of variables, the models did not perform as well (Table 2).

## DISCUSSION

One of us (JWA) first described the existence of *T. marmoratus* in the research area through a sporadic observation in March 1998. Our results confirm the presence of *T. marmoratus* in six populations outside its documented range in a pocket near Caldas da Rainha. This pocket of *T. marmoratus* is fully surrounded by populations of *T. pygmaeus*. Considering that (1) the distance of c. 10 km that separates it from the main *T. marmoratus* distribution exceeds the dispersal capability of large bodied newts (Arntzen & Wallis, 1991; Thiesmeier & Kupfer, 2000) and (2) that the species are locally strongly parapatric, we conclude that the pocket equals to an enclave.

The variable ORCH has a stronger effect than its fellow explanatory variables in models 5–9 (Table 2). It would be inappropriate though to extrapolate any model with ORCH over wider areas because in Portugal extensive fruit growing is particular to the Caldas da Rainha region.

In the habitat preference model, we assume that strongly preferred habitat is of high quality for the species (Railsback *et al.*, 2003). So, if a model has a good fit, it is usually assumed that the species/habitat system is in equilibrium and that the species distribution will only change if the environment around also changes. This ignores, however, the effects of life history and dispersal. Areas may be suitable but out-of-reach, due for ex-

ample to unsuitable habitat in between realized and prospective ranges. The current distribution of marbled newts in central Portugal is best explained by *T. pygmaeus* moving north from its previous range and superseding *T. marmoratus* that only persisted in areas with ecological conditions more suitable for the species. Because *T. marmoratus* and *T. pygmaeus* occasionally hybridize, this scenario can, in principle, be tested by looking for *T. marmoratus* 'genetic footprints' in areas that are hypothesized to have been inhabited by *T. marmoratus* in the past (Arntzen & Wallis, 1991). Similarly, we predict the absence of *T. pygmaeus* 'genetic footprints' inside the enclave.

There are several accounts of moving hybrid zones in, for example, butterflies (Mallet *et al.*, 1990; Blum, 2002; Dasmahapatra *et al.*, 2002), birds (Rohwer *et al.*, 2001; Pearson, 2000), crayfish (Perry *et al.*, 2001), lizards (Hillis & Simmons, 1986), and amphibians (Arntzen, 1978; Arntzen & Wallis, 1991; see also Barton & Hewitt (1985; 116–119). Most studies provide direct evidence of hybrid zone movement through the tracing of genetically interacting species distributions over time. Our study utilizes a single temporal window and provides compelling evidence for spatial change in a mosaic hybrid zone nevertheless, through the demonstration of an enclave. The distance between *T. marmoratus* in the enclave and the main distribution is minimally 6 km (the distance between populations 13 and 15) and maximally 15 km (the distance between populations 22 and 23). Although we cannot pertinently exclude the presence of long-distance dispersal, such a scenario is unlikely, given the absence of *T. marmoratus* or genetically mixed individuals in populations 13 and 15. Similarly, we cannot exclude the possibility that a human introduction is responsible for the enclave. There is, however, no tradition of newt husbandry in Portugal, and a deliberate or accidental release is improbable.

To infer the direction of the movement in a hybrid zone, it is equally possible to follow a direct or an indirect strategy. Direct demonstrations employ two or more temporally separated observations, either on position (e.g. Hillis & Simmons, 1986 in *Pholidobolus* lizards) or on shape of the cline that separates in this case connects the hybridizing species (e.g. Dasmahapatra *et al.*, 2002 in *Anartia* butterflies). An indirect way is to look at

**Table 2** List of *a priori* and *a posteriori* models used to determine the relationship between the distribution of *Triturus marmoratus* and *Triturus pygmaeus* in the Caldas da Rainha area, Portugal. The parameter values were obtained by logistic regression analysis of presence data in the area (by contrasting the presence of one species with the presence of the other). The log likelihood of the models ( $-2 \ln L$ ), number of parameters in each model ( $K$ ), and the number of data points ( $n$ ) were used to calculate the small-sample Akaike Information Criterion (AICc). The hypothesis behind each model is also listed. Lowest AICc value is highlighted in bold.

| Model               | $-2 \ln L$ | $K$ | $n$ | AICc         | Hypothesis                              | Model equation |       |       |       |       |       |       |       |       |       |       |       |
|---------------------|------------|-----|-----|--------------|---|----------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
|                     |            |     |     |              |   | ORCH           | RELI  | TJUL  | FROD  | ALTI  | PRET  | HARD  | HUMI  | NDVI  | TEMP  | SULP  | ACID  |
| <i>A priori</i>     |            |     |     |              |   |                |       |       |       |       |       |       |       |       |       |       |       |
| 1                   | 25.50      | 1   | 30  | 53.14        | Extent of orchard plantations           | 2.85           |       |       |       |       |       |       |       |       |       |       | -1.23 |
| 2                   | 36.87      | 1   | 30  | 75.88        | Relief                                  |                | -1.04 |       |       |       |       |       |       |       |       |       | -0.12 |
| 3                   | 40.24      | 1   | 30  | 82.62        | July temperature                        |                |       | -0.42 |       |       |       |       |       |       |       |       | 0.05  |
| 4                   | 41.23      | 1   | 30  | 84.60        | Number of frost days                    |                |       |       | -0.04 |       |       |       |       |       |       |       | 0.08  |
| 5                   | 24.85      | 2   | 30  | 54.15        | Combination of model 1 and model 2      | 2.57           | -0.61 |       |       |       |       |       |       |       |       |       | -1.24 |
| 6                   | 24.02      | 3   | 30  | 54.96        | Model 5 plus altitude                   | 2.78           | -0.33 |       |       | -0.83 |       |       |       |       |       |       | -1.49 |
| 7                   | 24.74      | 3   | 30  | 56.40        | Model 5 plus precipitation              | 2.50           | -0.56 |       |       |       | -0.22 |       |       |       |       |       | -1.21 |
| 8                   | 24.77      | 3   | 30  | 56.47        | Combination of model 4 and model 5      | 2.62           | -0.57 |       | -0.18 |       |       |       |       |       |       |       | -1.25 |
| 9                   | 20.72      | 3   | 30  | <b>48.36</b> | Combination of model 3 and model 5      | 5.10           | -0.40 | -2.10 |       |       |       |       |       |       |       |       | -3.30 |
| 10                  | 28.52      | 6   | 30  | 72.70        | from Arntzen (unpublished)              |                | -0.98 | -0.44 |       |       |       | 0.16  | -1.81 | -0.87 | -1.88 |       | -0.68 |
| <i>A posteriori</i> |            |     |     |              |   |                |       |       |       |       |       |       |       |       |       |       |       |
| 11                  | 28.73      | 5   | 30  | 69.97        | SMR model                               |                |       |       |       | -0.92 |       | -3.07 |       | -2.88 | 0.79  | -0.38 | -0.72 |
| 12                  | 27.06      | 6   | 30  | 69.76        | Model 5 with ORCH replaced by SMR model |                | -1.06 |       |       | -0.70 |       | -3.18 |       | -2.22 | 0.90  | -0.75 | -0.77 |
| 13                  | 26.12      | 7   | 30  | 71.32        | Model 7 with ORCH replaced by SMR model |                | -0.74 |       |       | 0.00  | -1.63 | -3.69 |       | -2.70 | 0.24  | -0.48 | -0.90 |
| 14                  | 26.80      | 7   | 30  | 72.70        | Model 8 with ORCH replaced by SMR model |                | -1.14 |       | 0.42  | -0.92 |       | -3.19 |       | -2.14 | 1.02  | -0.84 | -0.83 |
| 15                  | 22.82      | 7   | 30  | 64.75        | Model 9 with ORCH replaced by SMR model |                | -2.17 | -2.82 |       |       |       | -4.53 |       | -2.07 | 2.24  | -0.31 | -2.01 |

SMR, stepwise multiple linear regression; ORCH, orchard land coverage; RELI, relief; TJUL, mean July temperature; FROD, frost days; ALTI, altitude; PRET, mean annual precipitation; HARD, hardness of subterranean water; HUMI, humidity of the air; NDVI, vegetation index; TEMP, mean annual temperature; SULP, sulphate content of subterranean water; ACID, acidity of the soil (details in Teixeira *et al.* 2001 and Arntzen, 2006).

disequilibrium measures. Cruzan (2005) showed in the wide *Piriqueta caroliniana/viridis* (flowering plants from the family Turneracea) hybrid zone that *P. viridis* alleles were moving north. This was done by showing that the southern border of the hybrid zone presented relatively high levels of disequilibria, indicating recent gene flow from parental populations south of the hybrid zone. Interestingly, we observed significant levels of linkage disequilibrium in two *T. marmoratus* populations (populations 23 and 31) that are both within the dispersal range of *T. pygmaeus* populations. This strengthens the argument that *T. pygmaeus* advances into *T. marmoratus* territory, even though we were unable to demonstrate current *T. pygmaeus* presence.

As a further test to our explanation, we predict that the presence of alien *T. marmoratus* genes in *T. pygmaeus* exceeds that of the reverse condition, in a spatial pattern consistent with enclave formation.

Despite the range expansion of *T. pygmaeus* in its northern border, the situation in the south of Spain and Portugal is different. Due to desertification and an intensification of agricultural practices, *T. pygmaeus* is losing many breeding sites and has been classified as 'near threatened' (Arntzen *et al.* 2004). This is consistent with recent evidence that suggests that climate warming will not only increase the northern range of species but also decrease the southern one (Thomas *et al.*, 2006).

If *T. pygmaeus* continues its competitive advance, the *T. marmoratus* enclave would eventually disappear. As yet, we have no indication on the speed of the process. Monitoring the area would provide an additional test to the hypothesis of *T. pygmaeus* expansion and document the speed of the process. It is remarkable that *T. marmoratus* is not only losing out to its sister-species at the southern edge of its range, but also to the related species *T. cristatus* at its northern edge. The advance of *T. cristatus* at the expense of *T. marmoratus* was estimated to occur at a speed of c. 1 km per year (Arntzen & Wallis, 1991). If range replacement would proceed at this speed at either side of its range, *T. marmoratus* would be squeezed out in c. 500 years.

## ACKNOWLEDGEMENTS

We are grateful to Armando da Costa Pais from Direcção Regional da Agricultura do Alto Oeste for discussion on the relevance of agricultural data for our work. The study was carried under the license from the ICN (Instituto de Conservação da Natureza) in accordance to National law for capturing wild fauna and was financed by FCT (Fundação para a Ciência e Tecnologia) research project POCTI/34110/99.

## REFERENCES

- Anderson, E.C. & Thompson, E.A. (2002) A model-based method for identifying species hybrids using multilocus genetic data. *Genetics*, **160**, 1217–1229.
- Arntzen, J.W. (1978) Some hypotheses on postglacial migrations of the fire-bellied toad, *Bombina bombina* (Linnaeus) and the yellow-bellied toad, *Bombina variegata* (Linnaeus). *Journal of Biogeography*, **5**, 339–345.
- Arntzen, J.W. (2006) From descriptive to predictive distribution models: a working example with Iberian amphibians and reptiles. *Frontiers in Zoology*, **3**, 8.
- Arntzen, J.W., Jehle, R., Bosch, J., Miaud, C., Tejado, M., Lizana, M., Martínez-Solano, I., Salvador, A., García-París, M. & Gil, E.R. (2004) *Triturus marmoratus*. IUCN 2006. 2006 IUCN red list of threatened species. (www.iucnredlist.org).
- Arntzen, J.W. & Wallis, G.P. (1991) Restricted gene flow in a moving hybrid zone of the newts *Triturus cristatus* and *T. marmoratus* in western France. *Evolution*, **45**, 805–826.
- Arntzen, J.W. & Wallis, G.P. (1999) Geographic variation and taxonomy of crested newts (*Triturus cristatus* superspecies): morphological and mitochondrial data. *Contribution to Zoology*, **68**, 181–203.
- Barton, N.H. & Hewitt, G.M. (1985) Analysis of hybrid zones. *Annual Review of Ecology and Systematics*, **16**, 113–148.
- Bierne, N., Borsa, P., Daguin, C., Jollivet, D., Viard, F., Bonhomme, F. & David, P. (2003) Introgression patterns in the mosaic hybrid zone between *Mytilus edulis* and *M. galloprovincialis*. *Molecular Ecology*, **12**, 447–461.
- Blum, M.J. (2002) Rapid movement of a *Heliconius* hybrid zone: evidence for phase III of Wright's shifting balance theory? *Evolution*, **56**, 1992–1998.
- Bridle, J.R., Baird, S.J.E. & Butlin, R.K. (2001) Spatial structure and habitat variation in a grasshopper hybrid zone. *Evolution*, **55**, 1832–1843.
- Burnham, K.P. & Anderson, D.R. (2002) *Model selection and multimodel inference: a practical-theoretic approach*, 2nd edn. Springer, New York.
- Cruzan, M.B. (2005) Patterns of introgression across an expanding hybrid zone: analysing historical patterns of gene flow using nonequilibrium approaches. *New Phytologist*, **167**, 267–278.
- Dasmahapatra, K.K., Blum, M.J., Aiello, A., Hackwell, S., Davies, N., Bermingham, E.P. & Mallet, J. (2002) Inferences from a rapidly moving hybrid zone. *Evolution*, **56**, 741–753.
- Direcção Geral do Ambiente (1995) *Atlas do Ambiente. Versão 1.1*. Direcção Geral do Ambiente. Ministério do Ambiente e dos Recursos Naturais. Lisboa, Portugal.
- Espregueira Themudo, G. & Arntzen, J.W. (2007) Molecular identification of marbled newts and a justification of the *Triturus marmoratus* and *T. pygmaeus* species status. *Herpetological Journal*, **17**, (in press).
- Excoffier, L.G., Laval & Schneider, S. (2005) Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online*, **1**, 47–50.
- Goudet, J. (1995) Fstat version 1.2: a computer program to calculate F-statistics. *Journal of Heredity*, **86**, 485–486.
- Harrison, R.G. (1990) *Hybrid zones: windows on evolutionary processes*. Oxford University Press, New York.
- Hewitt, G.M. (1988) Hybrid zones-natural laboratories for evolutionary studies. *Trends in Ecology & Evolution*, **3**, 158–167.
- Hillis, D.M. & Simmons, J.E. (1986) Dynamic change of a zone of parapatry between two species of *Pholidobolus* (Sauria: Gymnophthalmidae). *Journal of Herpetology*, **20**, 85–87.
- Instituto Nacional de Estatística (1999) *Recenseamento Geral da Agricultura*. (Available from [http://www.inec.pt/prodserv/Rga/rga99\\_2.asp](http://www.inec.pt/prodserv/Rga/rga99_2.asp)).

- International Institute for Geo-Information Science and Earth Observation (ITC) (1997) *Ilwis 2.1 for Windows*. ITC, Enschede, the Netherlands.
- Jaeger, R.G. (1970) Potential extinction through competition between two species of terrestrial salamanders. *Evolution*, **24**, 632–642.
- Jaeger, R.G. (1971) Competitive exclusion as a factor influencing the distributions of two species of terrestrial salamanders. *Ecology*, **52**, 632–637.
- Jehle, R., Arntzen, J.W., Burke, T., Krupa, A.P. & Hodl, W. (2001) The annual number of breeding adults and the effective population size of syntopic newts (*Triturus cristatus*, *T. marmoratus*). *Molecular Ecology*, **10**, 839–850.
- Jehle, R., Wilson, G.A., Arntzen, J.W. & Burke, T. (2005) Contemporary gene flow and the spatio-temporal genetic structure of subdivided newt populations (*Triturus cristatus*, *T. marmoratus*). *Journal of Evolutionary Biology*, **18**, 619–628.
- Jiggins, C.D. & Mallet, J. (2000) Bimodal hybrid zones and speciation. *Trends in Ecology & Evolution*, **15**, 250–255.
- Key, K.H.L. (1981) Species, parapatry, and the Morabine grasshoppers. *Systematic Zoology*, **30**, 425–458.
- Mallet, J., Barton, N., Lamas, G., Santisteban, J., Muedas, M. & Eeley, H. (1990) Estimates of selection and gene flow from measures of cline width and linkage disequilibrium in *Heliconius* hybrid zones. *Genetics*, **124**, 921–936.
- Pearson, S.F. (2000) Behavioral asymmetries in a moving hybrid zone. *Behavioral Ecology*, **11**, 84–92.
- Perry, W.L., Feder, J.L., Dwyer, G. & Lodge, D.M. (2001) Hybrid zone dynamics and species replacement between *Orconectes* crayfishes in a northern Wisconsin lake. *Evolution*, **55**, 1153–1166.
- Pritchard, J.K., Stephens, M. & Donnelly, P. (2000) Inference of population structure using multilocus genotype data. *Genetics*, **155**, 945–959.
- Railsback, S.F., Stauffer, H.B. & Harvey, B.C. (2003) What can habitat preference models tell us? Tests using a virtual trout population. *Ecological Applications*, **13**, 1580–1594.
- Rand, D.M. & Harrison, R.G. (1989) Ecological genetics of a mosaic hybrid zone: mitochondrial, nuclear, and reproductive differentiation of crickets by soil type. *Evolution*, **43**, 432–449.
- Rohwer, S., Bermingham, E. & Wood, C. (2001) Plumage and mitochondrial DNA haplotype variation across a moving hybrid zone. *Evolution*, **55**, 405–422.
- Schoorl, J. & Zuiderwijk, A. (1981) Ecological isolation in *Triturus cristatus* and *Triturus marmoratus* (Amphibia: Salamandridae). *Amphibia-Reptilia*, **1**, 235–252.
- Serviço Climático Nacional (1974) *Atlas climatológico de Portugal*, SCN, Lisboa.
- Sites, J.W. Jr, Morando, M., Highton, R., Huber, F. & Jung, R.E. (2001) Phylogenetic relationships of the endangered shenandoah salamander (*Plethodon shenandoah*) and other salamanders of the *Plethodon cinereus* group (Caudata: Plethodontidae). *Journal of Herpetology*, **38**, 96–105.
- SPSS Inc. (2005) *SPSS for Windows, Rel. 14.0.0*. SPSS Inc., Chicago, Illinois.
- Teixeira, J., Ferrand, N. & Arntzen, J.W. (2001) Biogeography of the golden-striped salamander *Chioglossa lusitanica*: a field survey and spatial modelling approach. *Ecography*, **24**, 618–624.
- Thiesmeier, B. & Kupfer, A. (2000) *Der kammolch — ein wasserdrahe in gefahr*. Beiheft 1 der Zeitschrift für Feldherpetologie, Laurenti-Verlag, Berlin.
- Thomas, C.D., Franco, A.M.A. & Hill, J.K. (2006) Range retractions and extinctions in the face of climate warming. *Trends in Ecology & Evolution*, **21**, 415–416.
- Weir, B.S. & Cockerham, C.C. (1984) Estimating F-statistics for the analysis of population structure. *Evolution*, **38**, 1358–1370.

## SUPPLEMENTARY MATERIAL

The following supplementary material is available for this article:

**Table S1** Allele frequencies in XLS format.

**Table S2** Clustering assignment of 31 sample sites in XLS format.

This material is available as part of the online article from: <http://www.blackwell-synergy.com/doi/abs/10.1111/j.1472-4642.2007.00373.x>  
(This link will take you to the article abstract).

Please note: Blackwell publishing are not responsible for the content or functionality of any supplementary materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.